



KONGERIKET NORGE
The Kingdom of Norway

PCT/NO 00 / 00079

09/936165

REC'D 22 MAR 2000

WIPO PCT

#8

Bekreftelse på patentsøknad nr

Certification of patent application no

NO 00/00079

1999 1111

4

Det bekreftes herved at vedheftede dokument er nøyaktig utskrift/kopi av ovennevnte søknad, som opprinnelig inngitt 1999.03.08

It is hereby certified that the annexed document is a true copy of the above-mentioned application, as originally filed on 1999.03.08

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

2000.03.14

Freddy Strømmen

Freddy Strømmen
Seksjonsleder

Ellen B. Olsen

Ellen B. Olsen



PATENTSTYRET
Styret for det industrielle rettsvern

lc

PATENTSTYRET

08.MAR99 991111

Applicant: Norsk Hydro ASA
N-0240 OSLO

Attorney: Rita Lillegraven
Norsk Hydro ASA
N-0240 OSLO

Inventors: Kari Aasbø
Børsesjøveien 181
N-3719 SKIEN

Harald Breivik
Uranusveien 22
N-3942 SKJELSVIK

Title: "A new silage aid, process for preparing this silage aid and use of this silage aid.

P9923

This invention relates to a new silage aid, process for preparation of this silage aid and use of the silage aid.

When preparing silage from for instance fish waste, the raw material is treated with acid to obtain the optimum pH (3.5-4.5) with regard to enzymatic hydrolysis, and to prevent the growth of bacteria during storage.

During the silage process hydrolysis results in the formation of emulsions made up of an acidic water phase, an oil phase and solids. The degree of phase separation in these systems depends upon the type of raw material used. Oxygen is easily transferred through water, which unfortunately gives rise to oxidative degradation of the oil phase. This lipid oxidation results, of course, in a reduced nutritional value of the oil as an ingredient both for feed and for food. Likewise, it implies a reduced quality of the oleochemicals made from fish silage oil. To overcome these problems it would be a great advantage if an antioxidant well suited for the protection of fish oil could be present during the silage process.

The most favourable way to include an antioxidant would be as a constituent of the silage aid. Generally, it would be necessary to have at least 1% antioxidant in the silage aid. Water soluble antioxidants are easily included in an acidic silage aid. Unfortunately though, these compounds will not be able to protect the oil after separation since they will not be associated with the oil phase. Furthermore, the antioxidants that are food approved in Europe and/or the US at present, are mainly fat soluble compounds, *e.g.* 2,6-di-*tert*-butyl-4-methylphenol(BHT), 3-*tert*-butyl-4-hydroxyanisole(BHA), *tert*-butylhydroquinone(TBHQ), tocopherol and the gallates. All of these food approved antioxidants are also commercially available in formulations with emulsifying agents, but unfortunately the emulsifying agents are in general hydrophobic, and thus these formulations are not well suited to an acidic water medium.

The antioxidant ethoxyquin is an exception with regard to the properties described above. Ethoxyquin is a secondary, cyclic amine. Hence, it is well known that ethoxyquin is soluble in acids at low pH (<3), as this implies protonation of the amino group and thereby formation of a hydrophilic salt. As pH rises to 4-5 however, ethoxyquin will be

deprotonated and consequently, will again become fat soluble. Thus, by choosing the proper concentration of acid it is possible to have ethoxyquin in the form of a hydrophilic salt in the silage aid, while during the silage process the salt is deprotonated, becomes fat soluble and consequently will be able to protect the oil against lipid oxidation. Likewise, other antioxidants with an amino group may be suitable for this use. The preparation of ethoxyquin formate salt, and this salt dissolved in formic acid is disclosed in Norwegian patent application no. 851007.

Unfortunately though, ethoxyquin or other known antioxidants with an amino group, are not food approved. Therefore, when food approval has been required, no silage aid soluble antioxidants have been available to date.

Thus it is a main object of the present invention to provide a new silage aid comprising a food approved antioxidant.

This and other objects of the invention is achieved by the attached claims.

The invention will be further explained below.

The antioxidants BHT, BHA, tocopherol, TBHQ and propyl gallate(PG) are practically insoluble in water, and experiments have shown that their solubility in mineral acids (hydrogen chloride, sulphuric acid) is also quite low (see Table 1). Unexpectedly however, we have now found that these highly hydrophobic antioxidants are soluble in the short chain carboxylic acids formic, acetic and propionic acid (see Table 1). Further experiments have shown that when using a silage aid comprising BHA, TBHQ or PG dissolved in 85% formic acid, a superior quality of the fish oil product as compared to the product of the same process using only 85% formic acid (Table 2) is obtained. When the silage aid containing antioxidant is blended with the fish waste raw material, the hydrophobic antioxidant is associated with and protects the oil phase. Thus, we have found a method which is convenient on a technical scale and which leads to superior quality of the products of the silage process.

Generally, the amount of silage aid needed will depend upon the type of fish waste used and the choice of acid. Also, the amount of antioxidant needed, may depend upon the raw material used or the requirements regarding the stability of the isolated oil. This implies that the required amount of antioxidant dissolved in the acid may vary.

Furthermore, we have shown by experiments that the solubility of BHT in formic acid increases when BHA is present in the acid. This was a surprising result.

The silage aid might further comprise additives like, anti-microbiell compounds (e.g. ethyl benzoate or benzoic acid), anti-fungal compounds, anti-corrosive compounds, chelating compounds (e.g. citric acid), compounds improving the handling properties (e.g. glycerol), and oxygen scavengers.

The present invention also comprises to firstly dissolve the said antioxidants in a short chain carboxylic acid, and subsequently adding a mineral acid in the purpose of decreasing pH.

The short chain carboxylic acids according to this invention might be used either alone or in combination with their corresponding salts. Further, the aforementioned acids might be used as a mixture or as a mixture together with any of their salts.

The present invention is documented by experiments performed for a fish silage process. This invention will of course also prevail for other processes where acidic preservation is used. Fish silage shall only be considered as an example. The invention is applicable in acidic preservation of other organic by-products like slaughter waste, poultry waste and food waste, as well.

Table 1. Solubility (weight %) of antioxidants in different acids.

	BHA	TBHQ	PG	BHT	Toco- pherol
5 M hydrogen chloride ^a	< 0,1%	< 0,1%	< 0,1%	< 0,1%	< 0,1%
4 N sulphuric acid ^a	< 0,1%	< 0,1%	< 0,1%	< 0,1%	< 0,1%
85% formic acid ^b	>5%	>5%	>10%	< 0,5% ^c	< 0,5%
98-100% formic acid ^b	>10%	>10%	>10%	< 3%	< 0,5%
100% acetic acid ^b	>10%	>10%	>10%	>10%	>10%
100% propionic acid ^b	>10%	>10%	<10%	>10%	>10%

a) A mixture of antioxidant(40mg) and mineral acid(40g) was thoroughly shaken at 23°C. b) A mixture of the chosen amount of antioxidant(20-400mg) and acid(4g) was thoroughly shaken at 23°C. c) 0.25% BHT is not soluble in formic acid. When 0.25% BHA is added to the formic acid, BHT is soluble.

Example I: Experimental procedure for a lab-scale fish silage process.

The raw material, Atlantic salmon viscera, was ground in a kitchen grinder, and the resulting minced fish waste was thoroughly blended before it was portioned into separate batches each containing 500 gram. Silage aid (2% vol./wt.; 10mL) 85% formic acid with or without 0.75% wt./vol. of dissolved antioxidant was added, and the content of each batch was mixed to ensure a homogenous distribution. The batches was stored for eight days in an oven at 35±2°C, before a standard procedure for silage work up was conducted. This included warming the silage at 90±1°C for 1 minute, tempering, and finally separating the oil after centrifugation. The isolated oils were analysed to determine the POV and *p*-AV. Furthermore, weight gain (at 35±1°C) as a function of time was registered to determine the IP of the oils. The results are presented in Table 2.

Example II: Experimental procedure as in Example I, but the raw material in this series was whole herring. The results are presented in Table 2.

Table 2. Comparison of the results from analysis of various oil quality parameters^a, determined for fish silage oil produced^b using silage aid with or without dissolved antioxidant.

Example no.	Silage aid ^c	POV	<i>p</i> -AV	IP
I	Control; 85% HCOOH (formic acid)	3	24	1
	85% HCOOH incl. BHA (150ppm)	2	22	19
	85% HCOOH incl. TBHQ (150ppm)	1	9	62
II	Control; 85% HCOOH	29	29	0
	85% HCOOH incl. PG (150ppm)	7	19	4

a)POV(Peroxide value, Ph. Eur. V.3.4.5); determines the amount of primary oxidation products, hydroperoxides, in the oil. *p*-AV(*p*-Anisidine value, IUPAC 2.504); determines the amount of some secondary oxidation products, alkenals, in the oil. IP(induction period); a measure of the shelf life of the oil, given as the number of days it takes before the oil shows detectable weight gain due to oxygen absorption.

b)The experimental procedure for lab-scale silage experiments is given above in example I. c)The concentration of antioxidant given in parenthesis, is based on the amount of fish waste used in these experiments.



Claims

1. Silage aid comprising at least one antioxidant selected from the group 2,6-di-*tert*-butyl-4-methylphenol (BHT), 3-*tert*-butyl-4-hydroxyanisole (BHA), *tert*-butylhydroquinone (TBHQ), tocopherol and gallates; at least one short chain carboxylic acid; and optionally at least one salt of said acid.
2. Silage aid according to claim 1, wherein the short chain carboxylic acid is formic acid, acetic acid and/or propionic acid.
3. Silage aid according to claim 1 or 2, wherein the short chain carboxylic acid is formic acid of concentration 60-100%, preferred 83-98%.
4. Silage aid according to claim 1 or 2 claims, wherein the short chain carboxylic acid is acetic acid or propionic acid of concentration 60-100%, preferred 80-100%.
5. Silage aid according to any of the proceeding claims, wherein 0.1-10% of the antioxidant, preferred 0.3-2%, is dissolved in the short chain carboxylic acid.
6. Silage aid according to any of the proceeding claims, comprising an antioxidant selected from the group BHA, TBHQ and propyl galate (PG); and formic acid.
7. Silage aid according to any of the proceeding claims, comprising BHA and BHT; and formic acid.
8. Process for preparation of a silage aid comprising at least one antioxidant selected from the group 2,6-di-*tert*-butyl-4-methylphenol (BHT), 3-*tert*-butyl-4-hydroxyanisole (BHA), *tert*-butylhydroquinone (TBHQ), tocopherol and gallates; a short chain carboxylic acid; and optionally at least one salt of said acid, by dissolving the antioxidants in the acid.
9. Process according to claim 8, by dissolving 0.1-10% of the antioxidant, preferred 0.3-2%, in a short chain carboxylic acid of concentration 60-100%.

10. Process according to claim 8 or 9, by dissolving 0.1-10% of the antioxidant, preferred 0.3-2%, in formic acid, acetic acid and/or propionic acid of concentration 60-100%.


11. Process according to claim 8, 9 or 10, for preparation of a silage aid comprising 2,6-di-*tert*-butyl-4-methylphenol (BHT) and 3-*tert*-butyl-4-hydroxyanisole (BHA), and formic acid, by

a) dissolving BHA in the acid, and

b) subsequently, dissolving BHT in the solution obtained in step a).

12. Use of a silage aid comprising at least one antioxidant selected from the group 2,6-di-*tert*-butyl-4-methylphenol (BHT), 3-*tert*-butyl-4-hydroxyanisole (BHA), *tert*-butylhydroquinone (TBHQ), tocopherol and gallates; a short chain carboxylic acid; and optionally at least one salt of said acid, for protection of fish oil during a fish silage process.

13. Use of a silage aid comprising at least one antioxidant selected from the group 2,6-di-*tert*-butyl-4-methylphenol (BHT), 3-*tert*-butyl-4-hydroxyanisole (BHA), *tert*-butylhydroquinone (TBHQ), tocopherol and gallates; a short chain carboxylic acid; and optionally at least one salt of said acid, during preservation of organic by-products.



Abstract

This invention relates to a new silage aid, process for preparation of this silage aid and use of the silage aid. The silage aid comprises at least one antioxidant selected from the group 2,6-di-*tert*-butyl-4-methylphenol (BHT), 3-*tert*-butyl-4-hydroxyanisole (BHA), *tert*-butylhydroquinone (TBHQ), tocopherol and gallates; at least one short chain carboxylic acid; and optionally at least one salt of said acid. These antioxidants are food approved. The present silage aid is useful for preservation of organic by-products, and particularly for protection of fish oil during a fish silage process.

